

## WHAT IS CLAIMED IS:

1. An isolated or recombinant polynucleotide encoding an antigenic polypeptide comprising:
- 5 a) at least 17 contiguous amino acids from the mature polypeptide from SEQ ID NO: 2;
- b) at least 17 contiguous amino acids from the mature polypeptide from SEQ ID NO: 4;
- 10 c) at least 17 contiguous amino acids from the mature polypeptide from SEQ ID NO: 6;
- d) at least 17 contiguous amino acids from the mature polypeptide from SEQ ID NO: 8;
- e) at least 17 contiguous amino acids from the mature polypeptide from SEQ ID NO: 13;
- 15 f) at least 17 contiguous amino acids from the polypeptide from SEQ ID NO: 15;
- g) at least 17 contiguous amino acids from the polypeptide from SEQ ID NO: 17; or
- 20 h) at least 17 contiguous amino acids from the polypeptide from SEQ ID NO: 19.
2. The polynucleotide of Claim 1, encoding all of the polypeptide of:
- 25 a) mature SEQ ID NO: 2;
- b) mature SEQ ID NO: 4;
- c) mature SEQ ID NO: 6;
- d) mature SEQ ID NO: 8;
- e) mature SEQ ID NO: 13;
- f) SEQ ID NO: 15;
- 30 g) SEQ ID NO: 17; or
- h) SEQ ID NO: 19.
3. The polynucleotide of Claim 1, which hybridizes at 55° C, less than 500 mM salt, and 50% formamide to:
- 35 a) the mature polypeptide coding portion of SEQ ID NO: 1;

- 5 b) the mature polypeptide coding portion of SEQ ID NO: 3;
- c) the mature polypeptide coding portion of SEQ ID NO: 5;
- 10 d) the mature polypeptide coding portion of SEQ ID NO: 7;
- e) the mature polypeptide coding portion of SEQ ID NO: 12;
- f) the polypeptide coding portion of SEQ ID NO: 14;
- 15 g) the polypeptide coding portion of SEQ ID NO: 16; or
- h) the polypeptide coding portion of SEQ ID NO: 18.

4. The polynucleotide of Claim 3, comprising:

- 15 a) at least 35 contiguous nucleotides of the mature coding portion of SEQ ID NO: 1;
- b) at least 35 contiguous nucleotides of the mature coding portion of SEQ ID NO: 3;
- 20 c) at least 35 contiguous nucleotides of the mature coding portion of SEQ ID NO: 5;
- d) at least 35 contiguous nucleotides of the mature coding portion of SEQ ID NO: 7;
- e) at least 35 contiguous nucleotides of the mature coding portion of SEQ ID NO: 12;
- 25 f) at least 35 contiguous nucleotides of the coding portion of SEQ ID NO: 14;
- g) at least 35 contiguous nucleotides of the coding portion of SEQ ID NO: 16; or
- 30 h) at least 35 contiguous nucleotides of the coding portion of SEQ ID NO: 18.

5. An expression vector comprising the polynucleotide of Claim 1.

- 35 6. A host cell containing the expression vector of Claim 5, including a eukaryotic cell.

Sub B2  
CONT.

7. A method of making an antigenic polypeptide comprising expressing a recombinant polynucleotide of Claim 1.

5 8. A method for detecting a polynucleotide of Claim 1, comprising contacting said polynucleotide with a probe that hybridizes, under stringent conditions, to at least 25 contiguous nucleotides of:

- sub B2  
cont.
- 10 a) the polynucleotide comprising the signal processed coding portion of SEQ ID NO: 1;
- b) the polynucleotide comprising the signal processed coding portion of SEQ ID NO: 3;
- c) the polynucleotide comprising the signal processed coding portion of SEQ ID NO: 5;
- 15 d) the polynucleotide comprising the signal processed coding portion of SEQ ID NO: 7;
- e) the polynucleotide comprising the signal processed coding portion of SEQ ID NO: 12;
- f) the polynucleotide comprising the coding portion of SEQ ID NO: 14;
- 20 g) the polynucleotide comprising the coding portion of SEQ ID NO: 16; or
- h) the polynucleotide comprising the coding portion of SEQ ID NO: 18;

25 to form a duplex, wherein detection of said duplex indicates the presence of said polynucleotide.

9. A kit for the detection of a polynucleotide of Claim 1, comprising a compartment containing a probe that  
30 hybridizes, under stringent hybridization conditions, to at least 17 contiguous nucleotides of a polynucleotide of Claim 1 to form a duplex.

10. The kit of claim 9, wherein said probe is  
35 detectably labeled.

11. A binding compound comprising an antibody binding site which specifically binds to:

- 5 a) at least 17 contiguous amino acids from the signal processed form of SEQ ID NO: 2;
- b) at least 17 contiguous amino acids from the signal processed form of SEQ ID NO: 4;
- c) at least 17 contiguous amino acids from the signal processed form of SEQ ID NO: 6;
- 10 d) at least 17 contiguous amino acids from the signal processed form of SEQ ID NO: 8;
- e) at least 17 contiguous amino acids from the signal processed form of SEQ ID NO: 13;
- f) at least 17 contiguous amino acids from SEQ ID NO: 15;
- 15 g) at least 17 contiguous amino acids from SEQ ID NO: 17; or
- h) at least 17 contiguous amino acids from SEQ ID NO: 19.

20 12. The binding compound of Claim 11, wherein:

- a) said antibody binding site is:
- 25 1) selectively immunoreactive with a polypeptide of the signal processed form of SEQ ID NO: 2;
- 2) selectively immunoreactive with a polypeptide of the signal processed form of SEQ ID NO: 4;
- 30 3) selectively immunoreactive with a polypeptide of the signal processed form of SEQ ID NO: 6;
- 4) selectively immunoreactive with a polypeptide of the signal processed form of SEQ ID NO: 8;
- 35 5) selectively immunoreactive with a polypeptide of the signal processed form of SEQ ID NO: 13;

sub B2  
cont.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

5

- 6) selectively immunoreactive with a polypeptide of SEQ ID NO: 15;
- 7) selectively immunoreactive with a polypeptide of SEQ ID NO: 17;
- 8) selectively immunoreactive with a polypeptide of SEQ ID NO: 19; or

b) said binding compound is:

- 1) an antibody molecule;
- 2) a polyclonal antiserum;
- 3) detectably labeled;
- 4) sterile; or
- 5) in a buffered composition.

15

13. A method using the binding compound of Claim 11, comprising contacting said binding compound with a biological sample comprising an antigen, thereby forming a binding compound:antigen complex.

20

14. The method of Claim 13, wherein said biological sample is from a human, and wherein said binding compound is an antibody.

25

15. A detection kit comprising said binding compound of Claim 12, and:

- a) instructional material for the use of said binding compound for said detection; or
- b) a compartment providing segregation of said binding compound.

30

16. A substantially pure or isolated antigenic polypeptide, which binds to said binding composition of Claim 11, and further comprises at least 17 contiguous amino acids from:

35

- a) the signal processed polypeptide from SEQ ID NO: 2;
- b) the signal processed polypeptide from SEQ ID NO: 4;

sub B2  
cont 10

- 5
- sub B2  
cont.
- 10
- c) the signal processed polypeptide from SEQ ID NO: 6;
  - d) the signal processed polypeptide from SEQ ID NO: 8;
  - e) the signal processed polypeptide from SEQ ID NO: 13;
  - f) SEQ ID NO: 15;
  - g) SEQ ID NO: 17; or
  - h) SEQ ID NO: 19.

17. The polypeptide of Claim 16, which:

- 15
- 20
- 25
- 30
- a) comprises at least a fragment of at least 25 contiguous amino acid residues from a signal processed primate HDTEA84 protein;
  - b) comprises at least a fragment of at least 25 contiguous amino acid residues from a signal processed primate HSLJD37R protein;
  - c) comprises at least a fragment of at least 25 contiguous amino acid residues from a signal processed rodent RANKL protein; or
  - d) comprises at least a fragment of at least 25 contiguous amino acid residues from primate RANKL protein;
  - e) is a soluble polypeptide;
  - f) is detectably labeled;
  - g) is in a sterile composition;
  - h) is in a buffered composition;
  - i) binds to an sialic acid residue;
  - j) is recombinantly produced, or
  - k) has a naturally occurring polypeptide sequence.

18. The polypeptide of Claim 17, which:

- 35
- a) comprises at least 17 contiguous amino acids of the signal processed SEQ ID NO: 2;
  - b) comprises at least 17 contiguous amino acids of the signal processed SEQ ID NO: 4;

- c) comprises at least 17 contiguous amino acids of the signal processed SEQ ID NO: 6;
- d) comprises at least 17 contiguous amino acids of the signal processed SEQ ID NO: 8;
- e) comprises at least 17 contiguous amino acids of the signal processed SEQ ID NO: 13;
- f) comprises at least 17 contiguous amino acids of SEQ ID NO: 15;
- g) comprises at least 17 contiguous amino acids of SEQ ID NO: 17; or
- h) comprises at least 17 contiguous amino acids of SEQ ID NO: 19.

19. A method of modulating a precursor cell physiology or function comprising a step of contacting said cell with:

- a) a binding compound which binds to said polypeptide of Claim 16;
- b) an HDTEA84 polypeptide;
- c) an HSLJD37R polypeptide; or
- d) a RANKL polypeptide.

20. The method of Claim 19, wherein said contacting is in combination with a TNF family ligand, or an antagonist of said TNF family ligand.